

SEA ICE THICKNESS FROM KINEMATICS

Ron Kwok (1), Ben Holt (2)

(1) Jet Propulsion Laboratory
Mail Stop 300-235, 4800 Oak Grove Drive
Pasadena, 91109, CA
Email: ron@ampersand.jpl.nasa.gov

(2) Jet Propulsion Laboratory
Mail Stop 300-323, 4800 Oak Grove Drive
Pasadena, 91109, USA
Email: ben@pacific.jpl.nasa.gov

Abstract:

The RADARSAT synthetic aperture radar is currently providing repeat surveys of the Arctic Ocean sea ice on a near 3-day basis with its wide-swath ScanSAR mode. For the first time, we can produce basin-scale estimates of sea ice age and thickness from observations of ice motion derived from sequential SAR images of the Arctic Ocean. The direct observations we extract from the SAR imagery are the deformation of Lagrangian elements (cells) which are areas enclosed by polygons defined by vertices on the ice cover. The time-dependent deformation of these cells are computed from the motion of the vertices obtained by tracking these points in SAR imagery. From the record of deformation, we estimate the ice age and thickness within each of these cells. Each cell covers an initial area of approximately 10 km by 10 km. More than 60,000 of cells are used to cover the entire Arctic Ocean. These high-resolution observations give us an unprecedented view of the sea ice cover and are suitable for supporting polar investigations as well as operational applications.

We have produced estimates of sea ice deformation, thickness and age from two months of SAR maps of the Arctic Ocean from early November 1996 through January 1997. Here, we describe this dataset, their use for studies of the sea ice cover and the potential contribution of ENVISAT in helping to extend these observations into the future. These data products and a description of their format are available at a website at the Jet Propulsion Laboratory (<http://www-radar.jpl.nasa.gov/rgps>).